



Remdesivir

Per U.S. FDA, for treatment of suspected or confirmed COVID-19 patients

The Remdesivir PowerPoint is on the HPH COVID Intranet page for additional info.

To handle, **RN wears CHEMO gloves + CHEMO gown + 2nd CHEMO gloves.**

SUBMIT AN RL EVENT REPORT FOR ANY SUSPECTED ADVERSE EVENT!

Who can receive it?



Adult/peds w/ severe disease: SpO2 \leq 94% on RA, requiring supplemental O2, mechanical ventilation, or ECMO

Call MD if hypersensitivity presents in the form of:

↓ BP, ↓ HR, ↑ HR, SOB, wheezing, N/V, diaphoresis, shivering



Monitoring

- **Prior** - Check hepatic and renal fxn
- **During** - Telemetry in place, Monitor BP & HR initially, q15mins x 2, q30mins x 4. Call for SBP < 90 or HR < 60
- **Daily labs** - serum chemistries, CBC, ALT, AST, bili, and alkaline phosphatase




Remdesivir should not be administered simultaneously with any other medication.

There is no compatibility info available.

Infuse through secondary IV set

Post-admin flush:

- Ensure pt. receives all medication
- Volume of NS greater than the priming volume of the tubing



This med must be programmed to infuse over 120 mins using the Guardrails Drug Library & requires dual sign off for administration.