

PPE for DIRECT PATIENT CAREGIVERS for Hospitalized Patients: MD, RN, APRN, PA, CNA, CCT, RT, PT, OR, SLP, Radiology Techs, Clinical Pharmacists, Dietitians, etc.

DIRECT PATIENT CARE = Touching/examining patient or contact for more than 10 min and within 6 feet

For any rule out or confirmed COVID patient go to [[LINK](#)]

General Patients: No Known COVID Risk

1




surgical mask

AND

OR



goggles



face shield

Headgear remains on health care worker through multiple patients (i.e., do not replace per patient)


- 2 Fresh gloves for all patient contact. Gown for contact/droplet precaution.
- 3 Break back tie in order to don gown over head gear.
- 4 Proper hand hygiene.

Aerosolizing Procedure on ANY Patient



N-95

and



full face shield

1

OR



CAPR

Headgear remains on health care worker through multiple patients (i.e., do not replace per patient)

- 2 Fresh gloves and gown.
Break back tie in order to don gown over head gear.
- 3 Proper hand hygiene.

Use N-95 with face shield or CAPR if doing an aerosolizing procedure (i.e. intubation) or if patient is critically ill.

AEROSOLIZING PROCEDURES:

Nebulized Medication
Non-Invasive Ventilation
BiPAP/CPAP/High flow O₂
High Frequency Ventilation
Chest Physiotherapy
Manual Ventilation
Intubation
Extubation

Bronchoscopy/BAL
Laryngoscopy
Tracheostomy
PFTs
Open Suction
CPR
Second Stage of Labor
(Obstetric Patients)



CREATING A HEALTHIER HAWAII