

# Risk Tiers

	Tier 1 (Low Risk)	Tier 2 (Moderate Risk)	Tier 3 (High risk)
Patient Criteria	<ul style="list-style-type: none"> <li>&lt; 60 &amp; Healthy</li> </ul>	<ul style="list-style-type: none"> <li>60-75</li> <li>&lt;60 with any of the following: DM, asthma, COPD, smoker, HTN, CKD, pregnant</li> </ul>	<ul style="list-style-type: none"> <li>75 and greater</li> <li>&lt;75 with any of the following: CVD, ESRD, immuno-compromise, advanced COPD, cirrhosis, cancer, frailty or multiple morbid conditions</li> </ul>
	and	or	or
Symptoms	<ul style="list-style-type: none"> <li>Non-Specific, mild URI Sx or mild cough</li> </ul>	<ul style="list-style-type: none"> <li>More pronounced URI Sx or cough</li> <li>Fever</li> <li>SOB</li> </ul>	<ul style="list-style-type: none"> <li>Cough with pain, wheezing or shortness of breath lasting longer than 6 days; DOE, or chest tightness</li> </ul>
	and	or	or
Data (if available)	<ul style="list-style-type: none"> <li>Normal BP, RR, Pulse Ox and Pulse (mild tachycardia is normal if febrile)</li> <li>Normal Chest X-ray</li> </ul>	<ul style="list-style-type: none"> <li>Significant leuko/lymphopenia</li> <li>Elevated CRP</li> <li>Pulse Ox &lt;100% but &gt;93%</li> <li>Pulse &gt;100</li> </ul>	<ul style="list-style-type: none"> <li>Hypoxemia (93% or less)</li> <li>Infiltrates on CXR or CT Chest</li> </ul>
	and	or	or
Support System	<ul style="list-style-type: none"> <li>Able to self-isolate</li> <li>Adequate Support</li> </ul>	<ul style="list-style-type: none"> <li>Otherwise Tier 1 but <u>uncertain</u> support</li> </ul>	<ul style="list-style-type: none"> <li>Otherwise lower tier, but <u>unstable</u> support system</li> </ul>

# Risk Tiers

	Tier 1 (Low Risk)	Tier 2 (Moderate Risk)	Tier 3 (High risk)
Clinical Course	<ol style="list-style-type: none"> <li>1. Tier 1 with stable or improving symptoms</li> <li>2. <u>Previously Tier 2 patient</u> with improving symptoms over 6 days</li> </ol>	<ol style="list-style-type: none"> <li>1. Tier 2 with stable symptoms</li> <li>2. <u>Previously Tier 1 patient</u> with symptoms not improving over 6 days</li> <li>3. <u>Previously Tier 3 patient</u> with improving symptoms over 6 days</li> </ol>	<ol style="list-style-type: none"> <li>1. Tier 3 with stable symptoms</li> <li>2. <u>Previously Tier 2 patient</u> with symptoms not improving over 6 days</li> <li>3. <u>Previously Tier 1 patient</u> with worsening symptoms (clinical judgement.)</li> </ol>
When to escalate to next Tier	<p>With worsening of any symptoms - or - symptoms not improving over 6 days</p>	<p>With worsening of respiratory symptoms, pleuritic pain, hemoptysis, Pulse Ox &lt;93%, decreased urine output (&lt;3 voids in 24 hours), decreased fluid intake (&lt; 24 ounces of fluid in 24 hours), or signs of dehydration</p>	<p>With any worsening of symptoms:</p> <p>Contact: PCP, Urgent Care, or call 911</p>

