

PPE for Respiratory Evaluation Clinics

Patient



Place and keep surgical masks on patients except while examining.

All Patients with Respiratory Symptoms



surgical mask, face shield

Headgear remains on health care worker through multiple patients (i.e., do not replace per patient)

2 Fresh gloves for every patient.

Gown remains on health care worker unless soiled. Break back tie in order to don gown over head gear.


3 Proper hand hygiene.

Swabbers, Aerosolizing Procedures or Direct Patient Care

This includes staff who will perform tasks within patient care spaces for prolonged periods of time (≥ 15 minutes cumulative)



N-95



full face shield

1

OR



CAPR

Headgear remains on health care worker through multiple patients (i.e., do not replace per patient)

2

Fresh gloves between patients. Gown remains on health care worker unless soiled or doffing for breaks or lunch. Break back tie in order to don gown over head gear.

3

Proper hand hygiene.

Use N-95 with face shield or CAPR if doing an aerosolizing procedure (i.e. intubation) or if patient is critically ill.

AEROSOLIZING PROCEDURES:

Nebulized Medication
Non-Invasive Ventilation
BiPAP/CPAP/High flow O2
High Frequency Ventilation
Chest Physiotherapy
Manual Ventilation
Intubation
Extubation
Tracheostomy
PFTs
Open Suction

CPR
Second Stage of Labor (Obstetric Patients)
Endoscopic Procedure

HAWAII PACIFIC HEALTH | KAPI'OLANI PALI MOMI STRAUB WILCOX

CREATING A HEALTHIER HAWAII